







Table 3--Vegetables, selected: Fresh and processing per capita utilization, 1977-2001

Year	Asparagus			Broccoli			Carrots			Cauliflower			Total		
	Fresh	Canning	Freezing	Fresh	Freezing	Fresh	Canning	Freezing	Fresh	Freezing	Fresh	Canning	Freezing	Fresh	Freezing
-- Pounds, farm weight --															
1977	0.3	0.5	0.2	1.2	1.2	5.3	1.9	1.8	1.1	0.7	7.9	2.4	3.9		
1978	0.3	0.4	0.2	1.0	1.4	5.3	1.6	1.8	0.8	0.8	7.4	2.0	4.2		
1979	0.3	0.3	0.2	1.2	1.4	5.9	1.8	1.9	1.1	0.7	8.5	2.1	4.2		
1980	0.3	0.4	0.1	1.4	1.4	6.2	1.7	1.7	1.1	0.8	9.0	2.1	4.0		
1981	0.3	0.4	0.1	1.7	1.5	6.1	1.5	1.9	1.4	0.9	9.5	1.9	4.4		
1982	0.4	0.3	0.1	2.0	1.5	6.6	1.1	1.7	1.3	0.9	10.3	1.4	4.2		
1983	0.4	0.3	0.1	2.0	1.5	6.5	1.1	1.8	1.4	0.8	10.3	1.4	4.2		
1984	0.4	0.3	0.1	2.5	1.8	6.7	1.8	2.1	1.8	0.9	11.4	2.1	4.9		
1985	0.5	0.3	0.1	2.6	1.9	6.5	1.3	1.8	1.8	0.9	11.4	1.6	4.7		
1986	0.6	0.3	0.1	3.0	1.7	6.5	1.0	1.8	2.2	0.9	12.3	1.3	4.5		
1987	0.6	0.3	0.1	3.1	2.2	8.3	0.9	2.1	2.1	0.9	14.1	1.2	5.3		
1988	0.6	0.3	0.1	3.8	2.4	7.1	1.0	2.3	2.2	0.9	13.7	1.3	5.7		
1989	0.6	0.3	0.1	3.8	2.2	8.1	0.9	2.5	2.3	0.8	14.8	1.2	5.6		
1990	0.6	0.3	0.1	3.4	2.2	8.3	1.1	2.3	2.2	0.8	14.5	1.4	5.4		
1991	0.6	0.3	0.1	3.1	2.3	7.7	1.1	2.4	2.0	0.6	13.4	1.4	5.4		
1992	0.6	0.3	0.1	3.4	2.4	8.3	1.7	2.3	1.8	0.7	14.1	2.0	5.5		
1993	0.6	0.3	0.1	3.4	2.3	10.9	1.1	2.8	2.1	0.7	17.0	1.4	5.9		
1994	0.6	0.2	0.1	4.5	2.3	12.8	1.5	2.8	2.0	0.6	19.9	1.7	5.8		
1995	0.6	0.3	0.1	4.4	2.6	11.3	1.7	2.6	1.7	0.6	18.0	2.0	5.9		
1996	0.6	0.2	0.1	4.6	2.6	12.6	1.7	2.9	1.7	0.5	19.5	1.9	6.1		
1997	0.7	0.2	0.1	5.1	2.3	14.4	1.5	2.6	1.8	0.4	22.0	1.7	5.4		
1998	0.8	0.2	0.1	5.2	2.1	13.0	1.5	2.8	1.5	0.8	20.5	1.7	5.8		
1999	0.9	0.2	0.1	6.7	2.2	11.6	1.5	2.5	1.9	0.5	21.1	1.7	5.3		
2000	1.0	0.2	0.1	5.6	2.5	11.3	1.6	2.7	2.0	0.5	19.9	1.8	5.8		
2001 f	1.0	0.2	0.1	5.7	2.2	11.4	1.5	2.6	1.9	0.6	20.0	1.7	5.5		

f = ERS forecast.

Source: Economic Research Service, USDA.



Table 5--Potatoes: U.S. per capita utilization, by category, 1977-2001 1/

Year	Total	Fresh	Total	Freezing	Processing		
					Chips & shoe-strings	Dehydrating	Canning
-- Pounds, farm weight --							
1977	122.1	50.1	72.0	42.2	16.2	11.4	2.2
1978	119.5	46.0	73.5	42.6	16.5	12.1	2.3
1979	117.8	49.3	68.5	38.5	16.7	11.2	2.1
1980	114.7	51.1	63.6	35.4	16.5	9.8	1.9
1981	116.5	45.8	70.7	41.5	16.6	10.8	1.8
1982	115.0	47.1	67.9	38.6	17.0	10.4	1.9
1983	118.7	49.8	68.9	39.2	17.8	10.0	1.9
1984	122.1	48.3	73.8	43.7	18.0	10.3	1.8
1985	122.4	46.3	76.1	45.4	17.6	11.2	1.9
1986	125.9	48.8	77.1	46.3	18.1	10.9	1.8
1987	126.0	47.9	78.1	47.9	17.6	10.8	1.8
1988	122.3	49.6	72.7	43.3	17.1	10.4	1.9
1989	127.0	50.0	77.0	46.8	17.4	10.8	2.0
1990	124.1	46.8	77.3	46.5	16.4	12.6	1.8
1991	134.5	50.4	84.1	51.2	17.3	13.9	1.7
1992	130.6	48.6	82.0	50.1	17.2	12.9	1.8
1993	137.7	50.5	87.2	53.9	17.8	13.8	1.7
1994	138.3	50.2	88.1	56.3	16.7	13.4	1.7
1995	138.8	49.9	88.9	56.9	16.6	13.4	2.0
1996 r	147.2	50.7	96.5	61.1	16.7	16.9	1.8
1997 r	144.0	49.4	94.6	60.4	16.2	16.2	1.8
1998 r	141.0	47.9	93.1	59.1	15.1	17.2	1.5
1999 r	139.8	49.2	90.6	59.5	16.3	13.0	1.8
2000 p	147.9	51.1	96.8	60.4	17.0	17.5	1.9
2001 f	144.4	49.3	95.1	60.1	16.3	16.9	1.8

r = Revised. p = Preliminary. f = ERS forecast.

1/ Calendar-year estimates on a fresh-equivalent basis.

Source: Economic Research Service, USDA.

Table 6--Dry edible beans: U.S. per capita use for selected classes, calendar year 1977-2001

Year	Pinto	Navy	Great Northern	Red kidney	All limas	Other 1/	Total
-- Pounds per person --							
1977	2.0	2.1	0.4	0.5	0.3	1.1	6.4
1978	1.9	1.2	0.3	0.5	0.3	0.9	5.1
1979	2.0	2.0	0.4	0.5	0.3	1.2	6.4
1980	0.9	2.0	0.4	0.5	0.3	1.3	5.4
1981	1.9	1.2	0.3	0.3	0.3	1.4	5.4
1982	3.2	1.5	0.5	0.5	0.3	0.5	6.5
1983	2.4	1.6	0.5	0.5	0.3	1.2	6.5
1984	1.7	1.7	0.5	0.4	0.3	0.5	5.1
1985	2.8	1.8	0.5	0.4	0.4	1.2	7.1
1986	2.6	1.2	0.5	0.5	0.4	1.4	6.6
1987	2.2	1.4	0.2	0.4	0.2	0.8	5.2
1988	3.4	1.1	0.5	0.3	0.2	1.4	6.9
1989	2.2	1.1	0.4	0.4	0.3	1.5	5.9
1990	3.1	1.3	0.4	0.6	0.2	1.0	6.6
1991	3.3	1.5	0.5	0.5	0.2	1.3	7.3
1992	3.8	1.5	0.5	0.6	0.2	1.1	7.7
1993	3.6	1.2	0.4	0.6	0.2	1.4	7.4
1994	3.3	1.9	0.4	0.6	0.2	1.4	7.8
1995	3.3	1.7	0.4	0.6	0.2	1.5	7.7
1996	3.5	1.3	0.5	0.6	0.2	1.4	7.5
1997 r	3.6	1.4	0.4	0.6	0.2	1.4	7.5
1998 r	3.5	1.2	0.4	0.6	0.2	1.5	7.4
1999 r	3.6	1.2	0.4	0.6	0.2	1.6	7.6
2000 p	3.6	1.2	0.4	0.6	0.2	1.8	7.8
2001 f	3.5	1.1	0.5	0.6	0.2	1.6	7.5

p= Preliminary. r = revised. f = ERS forecast. 1/ Includes black, small red, pink, cranberry, garbanzo, blackeye, and all other dry edible beans.

Source: Economic Research Service, USDA.

Table 7--Mushrooms, all: U.S. per capita utilization, 1977-2001

Crop year	Fresh	Processing	Total
-- Pounds, farm weight --			
1977	0.86	1.62	2.50
1978	1.02	1.66	2.70
1979	1.13	1.73	2.80
1980	1.20	1.53	2.70
1981	1.38	1.53	2.90
1982	1.44	1.51	2.90
1983	1.64	1.81	3.40
1984	1.77	1.76	3.60
1985	1.78	1.81	3.60
1986	1.87	1.88	3.80
1987	1.92	1.64	3.50
1988	1.98	1.54	3.50
1989	2.04	1.53	3.50
1990	2.00	1.70	3.70
1991	1.93	1.75	3.68
1992	2.00	1.69	3.69
1993	1.99	1.71	3.70
1994	2.05	1.96	4.01
1995	2.05	1.76	3.81
1996	2.11	1.86	3.97
1997	2.35	1.68	4.03
1998	2.50	1.44	3.94
1999	2.54	1.62	4.16
2000 p	2.62	1.67	4.29
2001 f	2.64	1.66	4.30

p = Preliminary. f = ERS forecast.

Source: Economic Research Service, USDA.