



COVID-19 Working Paper: Obesity Prevalence Among U.S. Adult Subpopulations During the First Year of the COVID-19 Pandemic

Mariah D. Ehmke and Brandon J. Restrepo

This paper is published through USDA, Economic Research Service's (ERS) COVID-19 Working Paper series. This temporary Working Paper series is designed to publicly release preliminary analyses relevant to the impacts of the COVID-19 pandemic on agriculture, food, the environment, and rural America in a timely manner. ERS' COVID-19 Working Papers have not undergone the review and editorial process generally accorded official ERS publications, but ERS economists and social scientists reviewed them through an expedited process.

The findings and conclusions in this COVID-19 Working Paper are those of the authors and should not be construed to represent any official USDA or U.S. Government determination or policy.

Abstract

Data from the 2011–20 Behavioral Risk Factor Surveillance System were leveraged to estimate obesity disparities prior to the spring 2020 arrival of the Coronavirus (COVID-19) pandemic and during the first year of the pandemic, and then intra-pandemic changes in adult obesity rates were estimated across various population subgroups. Adult obesity was modeled as a function of various demographic and socioeconomic characteristics—household composition, urbanicity, Census region, age, race, gender, income, and education—over pre-pandemic and pandemic periods using a linear regression model estimated by ordinary least squares. The regression coefficients were then used to calculate the pre-pandemic and intra-pandemic obesity rates for the overall population and by subpopulation. Overall U.S. adult obesity prevalence was significantly higher during the first year of the pandemic by 1.3 percentage points (pp). This amounts to an obesity increase of 3.2 percent when compared with an obesity rate of 40.7 percent over the pre-pandemic period from January 1, 2019, to March 12, 2020. The net societal increase in early pandemic obesity rates was not evenly distributed across subpopulations. Obesity rates significantly increased by a larger amount for adults in subgroups with a lower accumulation of long-term human capital (such as education and skills) and greater potential need for institutional and societal support. Intra-pandemic obesity growth rates were significantly higher by 5.6 percent among adults with annual household incomes that qualified for benefits from the Supplemental Nutrition Assistance Program (SNAP) and by 7.3 percent for adults whose education level was below a high school diploma. Higher intra-pandemic growth rates for obesity were also observed for young adults (aged 20–39) (5.6 percent) and adults aged 20 years or older living in the West Census region and west to the Pacific Ocean, which includes all States from Montana, Wyoming, Colorado, and New Mexico (7.6 percent).

Keywords: COVID-19, Coronavirus, pandemic, obesity, rural obesity, urban obesity, household characteristics, race, ethnicity, gender, age, employment, income, education, Behavioral Risk Factor Surveillance System, BRFSS.

Acknowledgments

The authors thank the anonymous reviewers for technical peer reviews. They also thank Debbie Rubas, Jay Variyam, and Guijing Wang of USDA, Economic Research Service (ERS) for their comments, and Elaine Symanski, Jana Goldman, and Adele Wilcoxon of ERS for editorial and design services.

About the Authors

Mariah Ehmke and Brandon Restrepo are research agricultural economists in the Diet, Safety, and Health Economics Branch of the Food Economics Division of USDA, Economic Research Service.

Economic Research Service

www.ers.usda.gov

Use of commercial and trade names does not imply approval or constitute endorsement by USDA.

To ensure the quality of its research reports and satisfy governmentwide standards, ERS requires that all research reports with substantively new material be reviewed by qualified technical research peers. This technical peer review process, coordinated by ERS' Peer Review Coordinating Council, allows experts who possess the technical background, perspective, and expertise to provide an objective and meaningful assessment of the output's substantive content and clarity of communication during the publication's review.

In accordance with Federal civil rights law and U.S. Department of Agriculture (USDA) civil rights regulations and policies, the USDA, its Agencies, offices, and employees, and institutions participating in or administering USDA programs are prohibited from discriminating based on race, color, national origin, religion, sex, gender identity (including gender expression), sexual orientation, disability, age, marital status, family/parental status, income derived from a public assistance program, political beliefs, or reprisal or retaliation for prior civil rights activity, in any program or activity conducted or funded by USDA (not all bases apply to all programs). Remedies and complaint filing deadlines vary by program or incident.

Persons with disabilities who require alternative means of communication for program information (e.g., Braille, large print, audiotape, American Sign Language, etc.) should contact the responsible Agency or USDA's TARGET Center at (202) 720-2600 (voice and TTY) or contact USDA through the Federal Relay Service at (800) 877-8339. Additionally, program information may be made available in languages other than English.

To file a program discrimination complaint, complete the USDA Program Discrimination Complaint Form, AD-3027, found online at [How to File a Program Discrimination Complaint](#) and at any USDA office or write a letter addressed to USDA and provide in the letter all of the information requested in the form. To request a copy of the complaint form, call (866) 632-9992. Submit your completed form or letter to USDA by: (1) mail: U.S. Department of Agriculture, Office of the Assistant Secretary for Civil Rights, 1400 Independence Avenue, SW, Washington, D.C. 20250-9410; (2) fax: (202) 690-7442; or (3) email: program.intake@usda.gov.

USDA is an equal opportunity provider, employer, and lender.

Contents

Introduction	5
Background	6
Methods	11
Results	12
Summary Statistics	12
Main Findings	16
Overall Adult Population	16
Employment Status	17
Household Composition	18
Urbanicity/Rurality	19
Census Region	20
Age Group	21
Race and Ethnicity	22
Gender	23
Income Eligibility for SNAP	23
Educational Attainment	24
Discussion	25
Conclusion	27
References	29

What Is the Issue?

From March 2020 to June 2020, nearly 3 million U.S. residents had confirmed cases of COVID-19 and almost 130,000 nationwide had died of COVID-19 (Coronavirus Resource Center, 2022a, 2022b). Early in the pandemic, obesity was identified as a risk factor for medical complications and death from COVID-19 (Centers for Disease Control, 2022b; Sanchis-Gomar et al., 2020). The decades-long rise in adult obesity, from 30.5 percent in 1999–2000 to 42.4 percent in 2017–18, meant a higher percentage of adults faced increased risks from the COVID-19 virus. Public health measures to contain the pandemic altered behavioral, food access, and environmental constraints for millions of people living in the United States. Social pressures, including public health social distancing policies, food supply chain blockages, and workforce labor shortages, modified the day-to-day context of life. Many modifications may have worsened some obesity-related risk factors, such as sedentary behavior and snacking (Almandoz et al., 2022). Still, other pandemic-induced changes, like healthier at-home meal preparation, may have decreased obesity-related risk factors (Flanagan et al., 2020; Okrent and Zeballos, 2022; *The Economist*, 2020).

A nationally representative study shows a 3.0-percent increase in the rate of obesity in the overall adult U.S. population from spring 2019 to spring 2020 (Restrepo, 2022). Pre-pandemic obesity prevalence was not evenly distributed across society—adults in lower resource or marginalized situations tended to bear a higher proportion of the obesity burden (e.g., Befort et al., 2012; Grecu and Rothhoff, 2015). The different daily consequences of the pandemic for people living in the United States may also translate to uneven risks of obesity. As a result, pre-pandemic obesity disparities may have narrowed, remained constant, or widened across certain population subgroups (Belanger et al., 2020). This report measures changes in U.S. adult obesity rates overall and across a wide variety of demographic and socioeconomic subgroups during the first year of the pandemic—March 2020 to March 2021.

What Did the Study Find?

The first year of the COVID-19 pandemic compounded some pre-pandemic obesity prevalence disparities. The pandemic shock affected obesity rates differently across distinct population subgroups. While the U.S. obesity rate significantly increased from 40.7 percent to 42.0 percent ($p < 0.05$) or by 1.3 percentage points (pp) in the overall adult population during the first year of the pandemic, certain adult subpopulations experienced larger statistically significant increases in obesity rates. Adults whose annual household income was eligible for benefits from the Supplemental Nutrition Assistance Program (SNAP) had a 2.5 percentage point obesity increase during this time. Pronounced increases also occurred among younger adults aged 20–39 (2.0 pp), adults residing in the West Census region (2.8 pp) (Bureau of the Census, 2022), and adults with less than a high school diploma (3.3 pp). Only one subgroup examined in this study experienced a decline in obesity prevalence—adults living in the Northeast Census region—who reported a 1.1 pp decrease in obesity prevalence in the first year of the pandemic.

How Was the Study Conducted?

The data are from the 2011–20 annual waves of the Behavioral Risk Factor Surveillance System (BRFSS) from the Centers for Disease Control. After accounting for the BRFSS survey design, weighted BRFSS data are nationally representative of adults living in the United States (Centers for Disease Control, 2013). Self-reported Body Mass Index (BMI) was used to classify adults 20 years or older as having obesity ($BMI \geq 30$). BMI was corrected for self-reporting bias with data from the 2011–2012 to 2019–March 2020 cycles of the National Health and Nutrition Examination Survey (NHANES). The period from January 1, 2019, to March 12, 2020, is the baseline pre-pandemic period upon which intra-pandemic shifts in obesity rates were estimated.

Introduction

As COVID-19 emerged in the spring of 2020, it was soon known that obese individuals had a higher risk of severe adverse health outcomes, including death, from a COVID-19 infection than non-obese individuals (Centers for Disease Control, 2022b; Onyeaka et al., 2021; Sanchis-Gomar et al., 2020; Stefan et al., 2021). From March 2020 to June 2020, nearly 3 million U.S. residents had confirmed cases of COVID-19 and almost 130,000 had died of the disease (Coronavirus Resource Center, 2022a, 2022b). To prevent widespread death among such individuals and other at-risk U.S. residents (including elderly and immunocompromised individuals), local, State, and national authorities implemented public health measures (e.g., social distancing, schooling-at-home, and business closures) to limit the spread of infections and COVID-19 deaths beginning in March 2020 (Onyeaka et al., 2021).

These measures modified the day-to-day context of life in the United States (Centers for Disease Control, 2022a; Matthews et al., 2022). For some, daily modifications such as sedentary behavior and social isolation may have worsened obesity-related risk factors (Flanagan et al., 2020). At the same time, some adults reported new lifestyle activities, such as healthy at-home meal preparation and daily exercise. These activities may have decreased the risk of obesity among such adults (Flanagan et al., 2020; Okrent and Zeballos, 2022; Onyeaka et al., 2021; *The Economist*, 2020). However, overall adult obesity rates grew higher in the United States by the end of the first year of the COVID-19 pandemic (Restrepo, 2022).

The question remains whether the overall increase in adult obesity accelerated obesity-related health disparities during this time. Pre-pandemic obesity prevalence and growth rate differences arose across social, demographic, and economically defined groups (e.g., Hales et al., 2020a; Ogden et al., 2017). Certain population subgroups defined by employment status, household composition, urbanicity, Census region, age, race, gender, income, and education were more likely to suffer from obesity than the general population (e.g., Chen et al., 2018; Fryar et al., 2020; Ogden et al., 2017).

The authors measured changes in U.S. adult obesity rates across demographic and socioeconomic subgroups during the first year of the COVID-19 pandemic—March 2020 to March 2021. They analyzed pooled cross-sectional data from the 2011–20 Behavioral Risk Factor Surveillance System (BRFSS) to measure obesity rate changes. The nationally representative estimates were derived from a weighted cross-section of U.S. adults with data from before and during the first year of the pandemic (National Center for Chronic Disease Prevention and Health Promotion, 2022a). The data were analyzed using linear probability models to estimate changes in obesity rates across a wide variety of adult subpopulations.

The first year of the pandemic compounded some pre-existing obesity prevalence disparities. The pandemic shock affected obesity rates differently across distinct population subgroups. The authors found that large and statistically significant disparities arose across a wide variety of subgroups, including vulnerable segments of the population such as adults in households with children, those with annual household incomes that qualify for SNAP benefits, and those without a high school diploma.

This report serves to inform USDA's efforts to support food and nutrition security, which are essential elements to reduce the risk of obesity (U. S. Department of Agriculture, 2022). The results support efforts to improve nutritional education and counseling to reduce obesity and improve health equity (The White House, 2022).

In 2020, the COVID-19 pandemic arrived amidst sustained increases in obesity prevalence in the United States (Sanchis-Gomar et al., 2020). The pandemic induced unprecedented changes to community, household, and individual life from March 2020 to March 2021 (Centers for Disease Control, 2022a; Onyeaka et al., 2021; Thompson et al., 2020). Early on, public health researchers identified obesity as a risk factor for suffering from extensive illness and mortality from COVID-19 (Centers for Disease Control, 2022b; Sanchis-Gomar et al., 2020). As the pandemic evolved, varying obesity outcomes across the U.S. population were reported. Using data from emergency room and ambulatory care visits for individuals pre- and post-pandemic, Freedman et al. (2022) found a small average increase in weight (0.1 kg, or 0.2 lbs.) during the first year of the pandemic. However, researchers noted their sample was not nationally representative of the U.S. population. Restrepo (2022) used nationally representative Behavioral Risk Factor Surveillance System (BRFSS) data or self-reported cross-sectional surveillance data from the first year of the pandemic and found that the proportion of obese adults in the United States rose by 1.1 pp or 3.0 percent. These findings aligned with other research reports of rising adult obesity levels amidst the pandemic (Almandoz et al., 2022).

These discrepancies in findings suggest overall outcomes may not be representative of subgroup disparity trends. Within the population, certain groups may have had increased exposure to obesity risk that outweighed reductions in weight gain for other groups. For example, while Freedman et al. (2022) found a small average increase in weight across the population, they also found that women gained more weight than men and that young adults gained more weight relative to older adults. These findings suggest overall pandemic weight changes were not equally distributed across subpopulations. Such findings were consistent with long-term trends suggesting the obesity burden is not equally borne by U.S. residents.


In studies conducted prior to the pandemic, researchers found obesity is not evenly distributed across demographic, socioeconomic, and geographic segments of U.S. adults (Hales et al., 2020b; Ogden et al., 2017). Previous research on adult obesity-related health disparities points to a need to understand the possible differences in experiences across several demographic, socioeconomic, and geographic dimensions. The social determinants of health framework offers one approach to identify the most important demographic and socioeconomic characteristics affecting health status, including physical health as measured by obesity (Solar and Irwin, 2010) (see box, “Social Determinants of Health Framework and Heterogeneous Risks of Obesity Across Adult Subpopulations”).

Main Findings

Overall Adult Population

Figure 3 shows that the share of U.S. adults aged 20 and older with obesity significantly increased ($p < 0.05$) from 40.7 percent in the baseline pre-pandemic period to 42.0 percent during the first year of the COVID-19 pandemic (from March 13, 2020, to March 18, 2021).⁵ This statistically significant increase of 1.3 pp is not trivial when divided into the baseline pre-pandemic period obesity rate ($1.3/40.7 = 3.2$ percent).⁶

Figure 3
Obesity prevalence in the overall U.S. adult population, before and during the COVID-19 pandemic



Note: Statistical significance is indicated by ** $p < 0.05$.

Asterisks indicate the level of significance of the difference between the baseline pandemic period and the first year of the pandemic.

Source: USDA, Economic Research Service based on data collected by Behavioral Risk Factor Surveillance System.


Relative to the obesity rate in the baseline pre-pandemic period, the obesity rate was statistically indistinguishable over the first 3 months of the pandemic (March 2020 to May 2020) (40.8 percent) (figure 4). However, relative to the baseline pre-pandemic period, there were statistically significant increases in future pandemic periods ($p < 0.05$). Over the June 2020 to August 2020 period, approximately 3 months following the imposition of the initial pandemic restrictions, obesity prevalence significantly increased by 1.5 pp to 42.2 percent. Obesity prevalence remained at this elevated level, slightly increasing to 42.5 percent or 1.8 pp higher than the baseline pre-pandemic period from September 2020 to November 2020 and held steady at that level from December 2020 to March 2021.

⁵ In this analysis, the first year of the pandemic spans from March 13, 2020, to March 18, 2021.

⁶ This estimate slightly differs from Restrepo (2022) because that study used data from all U.S. areas, including U.S. territories, whereas this working paper uses data from the 50 U.S. States and Washington, D.C.

Figure 4

Obesity prevalence in the overall U.S. adult population, before and during stages of the COVID-19 pandemic



Note: Statistical significance is indicated by ** $p < 0.05$.

Asterisks indicate the level of significance of the difference between the baseline pandemic period and the given pandemic period.

Source: USDA, Economic Research Service based on data collected by Behavioral Risk Factor Surveillance System.


The above-mentioned changes are for the overall adult population. However, given the heterogeneity in obesity rates before the pandemic, varying obesity growth rates across different demographic characteristics, socioeconomic status, and geography may be expected. Therefore, the authors used the variables listed in table 1 to break down changes in obesity prevalence across a wide variety of adult subpopulations.

Employment Status

Prior to the pandemic, relative to employed respondents, obesity prevalence was slightly higher among those not employed (e.g., retired, unemployed, or stay-at-home parents) (figure 5). The obesity rate of employed respondents significantly increased by 1.2 pp ($p < 0.01$) or from a baseline pre-pandemic prevalence of 40.1 percent to 41.3 percent during the pandemic. The obesity rate of respondents who were not employed also increased by 1.2 pp, from 41.7 percent to 42.9 percent, but the change was not statistically significant.

Figure 5

Obesity prevalence in the U.S. by employment status, before and during the COVID-19 pandemic



Note: Statistical significance is indicated by *** $p < 0.01$.

Asterisks indicate the level of significance of the difference between the baseline pandemic period and the first year of the pandemic.


Source: USDA, Economic Research Service based on data collected by Behavioral Risk Factor Surveillance System.

Household Composition

Obesity rate changes also varied across groups according to marital status and whether there were children under 18 years of age in the household. Unmarried respondents without children reported a statistically insignificant obesity rate increase of 1 pp from 37.8 percent prior to the pandemic to 38.8 percent from March 2020 to March 2021 (figure 6). By contrast, single respondents with children reported an increase in their obesity rate that was almost twice as large (1.9 pp), which was statistically significant ($p < 0.05$). Married respondents without children reported one of the smallest obesity rate changes, with their obesity rate insignificantly rising from 41.2 percent to 41.8 percent. Conversely, married respondents with children reported a statistically significant increase ($p < 0.10$) in their obesity rate from 42.0 percent to 43.8 percent or 1.8 pp.

Figure 6

Obesity prevalence in the U.S. by household composition, before and during the COVID-19 pandemic



Note: Statistical significance is indicated by * $p < 0.10$ or ** $p < 0.05$.

Asterisks indicate the level of significance of the difference between the baseline pandemic period and the first year of the pandemic.


Source: USDA, Economic Research Service based on data collected by Behavioral Risk Factor Surveillance System.

Urbanicity/Rurality

Information on whether the respondent resided in an urban or rural area is available only from 2018 and later. Despite the smaller sample sizes by urbanicity, the authors were still able to estimate the baseline pre-pandemic period (2019 to pre-pandemic 2020) to the pandemic period change (figure 7).

Rural respondents' reported obesity rate was higher before the pandemic (46.9 percent) and held steady at that level during the first pandemic year. Urban respondents, however, started with a lower pre-pandemic obesity rate of 39.8 percent, which rose to 41.0 percent during the first year of the pandemic. The overall 1.2 pp rise in obesity rates among urbanites was statistically significant ($p < 0.05$).

Figure 7

Obesity prevalence in the U.S. by urbanicity/rurality before and during the COVID-19 pandemic

Note: Statistical significance is indicated by ** $p < 0.05$.

Asterisks indicate the level of significance of the difference between the baseline pandemic period and the first year of the pandemic.

Source: USDA, Economic Research Service based on data collected by Behavioral Risk Factor Surveillance System.


Census Region

The patterns by Census region were heterogeneous (figure 8). Interestingly, residents of the generally urbanized Northeast census region reported a small drop in obesity prevalence during the pandemic.⁷ Prior to the pandemic, the Northeast region respondents' obesity rate was 38.5 percent. From March 2020 to March 2021, the analysis indicates a 1.1 pp decrease in obesity prevalence for this group (37.4 percent). This change was statistically significant ($p < 0.01$). Respondents in all other geographic regions reported statistically significant obesity rate increases. The largest obesity rate increase was in the West, which is also generally urbanized, where the obesity rate across respondents increased from 36.7 percent to 39.5 percent ($p < 0.05$) or an increase of 2.8 pp. The two regions with the highest baseline pre-pandemic obesity rates were the Midwest and South. The baseline pre-pandemic obesity rate in the Midwest was 43.0 percent and 43.1 percent in the South. In the Midwest, respondents' reported weight increases translated into a 1.6 pp increase in the obesity rate during the pandemic (44.6 percent) ($p < 0.01$). This increase is about twice the size of the increase experienced in the South. There, respondents reported a 0.9 pp increase in the obesity rate from 43.1 to 44.0 percent ($p < 0.01$).

⁷ In our analytical sample with information on urbanicity/rurality (2018 and beyond), the share of respondents living in rural counties is 3.2 percent in the Northeast, 10.1 percent in the Midwest, 8.2 percent in the South, and 3.1 percent in the West.

Figure 8

Obesity prevalence by U.S. census region, before and during the COVID-19 pandemic



Note: Statistical significance is indicated by **p<0.05 or ***p<0.01.

Asterisks indicate the level of significance of the difference between the baseline pandemic period and the first year of the pandemic.


Source: USDA, Economic Research Service based on data collected by Behavioral Risk Factor Surveillance System.

Age Group

Figure 9 shows that the youngest adult age group examined had the largest obesity rate increase and was statistically significant. Relative to the baseline pre-pandemic period, obesity prevalence significantly increased by 2 pp among respondents 20 to 39 years old during the COVID-19 pandemic. This increase is over 1.5 times larger than the significant increase of 1.2 pp among respondents aged 40 to 59 years. The change in the obesity rate for people aged 60 years and older was even smaller and not statistically significant.

Figure 9

Obesity prevalence in the U.S. by age group before and during the COVID-19 pandemic



Note: Statistical significance is indicated by *p<0.10 or ***p<0.01. Asterisks indicate the level of significance of the difference between the baseline pandemic period and the first year of the pandemic.

Source: USDA, Economic Research Service based on data collected by Behavioral Risk Factor Surveillance System.

Race and Ethnicity


The variability in baseline obesity rates across racial and ethnic groups is very noticeable in figure 10. Prior to the pandemic, authors' estimates indicate that non-Hispanic Black respondents had the highest obesity rate at 47.9 percent or 7.2 pp above the full sample baseline pre-pandemic rate of 40.7 percent. American Indians, Alaskan Natives, and Native Hawaiian respondents' obesity rate of 47.6 percent was 6.9 pp above the full sample baseline pre-pandemic rate. Hispanic respondents' obesity rate was 46.6 percent or 5.9 pp higher than the full sample baseline pre-pandemic rate.

The non-Hispanic Asian subgroup was the racial minority with the lowest obesity rate prior to the pandemic. According to the survey respondent responses, the obesity rate of this group was 15.7 percent before the pandemic. The next lowest obesity rate, the non-Hispanic White subgroup, was nearly 3 times that of non-Hispanic Asians at 39.9 percent. This was just below the obesity rate of those of non-Hispanic Other races (40.0 percent).

After the first year of the pandemic, some racial groups' obesity rates were significantly higher. Specifically, non-Hispanic Black and White respondents reported statistically significant higher obesity rates relative to the baseline pre-pandemic period. Among non-Hispanic Black respondents, the obesity rate increased from 47.9 percent in the baseline pre-pandemic period to 49.5 percent in the first year of the pandemic, a statistically significant rise of 1.6 pp ($p < 0.10$). There was also a statistically significant 0.7 pp rise in non-Hispanic White participants' obesity rate. Among these respondents, the obesity rate increased significantly ($p < 0.05$) from 39.9 percent to 40.6 percent. Other racial and ethnic subgroups did not show statistically significant changes in their obesity rate, which may, in part, be due to their smaller sample sizes.

Figure 10

Obesity prevalence in the U.S. by race/ethnicity before and during the COVID-19 pandemic



AI/AN = American Indian or Alaskan Native and NH/PI = Native Hawaiian or other Pacific Islander.

Note: Statistical significance is indicated by * $p < 0.10$ or ** $p < 0.05$.

Asterisks indicate the level of significance of the difference between the baseline pandemic period and the first year of the pandemic.


Source: USDA, Economic Research Service based on data collected by Behavioral Risk Factor Surveillance System.

Gender

Obesity rates significantly increased among women during the pandemic compared to the baseline pre-pandemic period (figure 11). The obesity rate of women increased from 43.5 percent to 44.6 percent or by 1.1 pp ($p < 0.05$). Although the obesity rate increase among men was slightly larger at 1.3 pp, the estimate fell just outside the threshold of statistical significance ($p = 0.110$).

Figure 11

Obesity prevalence in the U.S. by gender before and during the COVID-19 pandemic



Note: Statistical significance is indicated by ** $p < 0.05$.

Asterisks indicate the level of significance of the difference between the baseline pandemic period and the first year of the pandemic.


Source: USDA, Economic Research Service based on data collected by Behavioral Risk Factor Surveillance System.

Income Eligibility for SNAP

Given that respondents came from households that varied by household size, estimates by annual household income alone are challenging to interpret. The authors used information on annual household income categories and household size in combination with poverty guidelines to determine whether respondents' annual household income made them eligible for SNAP. Three interesting patterns emerge. First, respondents whose income was not eligible for SNAP benefits had much lower baseline pre-pandemic obesity rates than respondents who were income eligible (figure 12). Second, while there was a 0.8 pp increase in the obesity rate among respondents whose income made them ineligible for SNAP benefits, the increase was not statistically significant. Third, by contrast, the increase in the obesity rate among respondents who were income-eligible for SNAP benefits was over twice the size (2.5 pp) and statistically significant ($p < 0.05$).

Figure 12

Obesity prevalence in the U.S. by SNAP (income) eligibility, before and during the COVID-19 pandemic



Note: Statistical significance is indicated by **p<0.05.

Asterisks indicate the level of significance of the difference between the baseline pandemic period and the first year of the pandemic.


Source: USDA, Economic Research Service based on data collected by Behavioral Risk Factor Surveillance System.

Educational Attainment

Across educational subgroups, the low education subgroup (those who attained below a high school diploma) reported the highest obesity rate increase (figure 13). Their pre-pandemic obesity rate was 45.1 percent. This rose by 3.3 pp or to an obesity rate of 48.4 percent during March 2020 to March 2021 ($p<0.01$). The educational subgroup with some college education also had a significant 1.5 pp ($p<0.10$) increase in their obesity rate from 42.8 percent to 44.3 percent over the same periods. Other educational subgroup respondents did not show a significant change in obesity rates.

Figure 13

Obesity prevalence in the U.S. by education attainment, before and during the COVID-19 pandemic



HS = high school.

Note: Statistical significance is indicated by * $p < 0.10$ or *** $p < 0.01$.

Asterisks indicate the level of significance of the difference between the baseline pandemic period and the first year of the pandemic.

Source: USDA, Economic Research Service based on data collected by Behavioral Risk Factor Surveillance System.

Discussion

Given the health risks associated with obesity (National Center for Chronic Disease Prevention and Health Promotion, 2022b), every statistically significant increase in obesity within a subgroup during the pandemic is concerning. The authors performed seemingly unrelated regressions (SUR) to compare the estimated change in the prevalence of obesity in the overall adult population—an increase of 1.3 pp—to each of the estimated obesity rate changes among each of the previously analyzed subgroups. After the SUR estimations, Wald tests of equality of regression coefficients indicated that the following subgroups exhibited obesity rate changes that were larger as well as statistically distinguishable from the obesity rate change in the overall adult population, with p-values under 0.10:

- Young adults aged 20–39 (estimated intra-pandemic increase = 2 pp)
- Adults living in the West Census region (estimated intra-pandemic increase = 2.8 pp)
- Adults in households with an annual income that was eligible for SNAP benefits (estimated intra-pandemic increase = 2.5 pp)
- Adults with below a high school diploma (estimated intra-pandemic increase = 3.3 pp)


One way to get a sense of the size of these increases is to compare them to average year-over-year (YOY) changes from 2011 to 2019. Using the full adult sample, the average pre-pandemic YOY change was 0.6 pp, so the intra-pandemic increase of 1.3 pp in the overall adult population was over twice as large as the earlier overall average YOY change. Homing in on the adult subgroups listed above, the subgroup-specific average pre-pandemic YOY changes are as follows:

- Young adults aged 20–39 (average pre-pandemic YOY change = 0.5 pp)
 - Estimated intra-pandemic increase was 2.5 or about 4 times larger
- Adults living in the West Census region (average pre-pandemic YOY change = 0.6 pp)
 - Estimated intra-pandemic increase was 2.8 or about 5 times larger
- Adults in households with an annual income that was eligible for SNAP benefits (average pre-pandemic YOY change = 1.3 pp)
 - Estimated intra-pandemic increase was 2.5 or about 2 times larger
- Adults with below a high school diploma (average pre-pandemic YOY change = 1 pp)
 - Estimated intra-pandemic increase was 3.3 or about 3 times larger

Therefore, the increases from the baseline pre-pandemic period to the first year of the COVID-19 pandemic were significant, ranging from about two times the size to about five times the size of the earlier average YOY changes.

In figure 14, each of the statistically significant increases ($p < 0.10$) is expressed as a share of each subgroup's baseline pre-pandemic obesity rate, and the increases are then sorted by baseline pre-pandemic obesity rate in ascending order. Young adults aged 20–39 had the smallest baseline pre-pandemic obesity rate (35.7 percent), and adults with below a high school diploma had the largest baseline pre-pandemic obesity rate (45.1 percent).

Figure 14
Estimated percent changes in obesity in the U.S. during the COVID-19 pandemic



SNAP = Supplemental Nutrition Program.

Note: Expressed as shares of pre-pandemic obesity rates, ranked from least to greatest pre-pandemic prevalence. Statistical significance is indicated by * $p < 0.10$, ** $p < 0.05$, or *** $p < 0.01$. Asterisks indicate the level of significance of the difference between that subgroup and overall adult population from a Wald test of equality of regression coefficients.

Source: USDA, Economic Research Service based on data collected by Behavioral Risk Factor Surveillance System.

Although young adults aged 20 to 39 years old started from a relatively low pre-pandemic rate of obesity, the prevalence of obesity increased by a substantial amount, which may, in turn, raise the risk of obesity and obesity-linked health effects in later adulthood (Poobalan and Aucott, 2016). These young adults reported a 2 pp intra-pandemic increase in obesity prevalence, or a 5.6-percent increase relative to the baseline pre-pandemic level. Most of these young adults are in their prime working years and have a stronger attachment to the labor force than older adults (Bureau of Labor Statistics, 2020). Compared with employed people who have a healthy weight, those with obesity are significantly more likely to miss work due to injuries or illnesses (Cawley et al., 2021).

In the baseline pre-pandemic period, those living in the West Census region had a lower prevalence of obesity than the three other Census regions (figure 8). However, during the first year of the pandemic, obesity prevalence in the West increased by 2.8 pp or 7.6 percent higher than the rate during the baseline pre-pandemic period. Together with a drop in obesity prevalence in the Northeast Census region, the West Census region's rank in obesity prevalence fell from the lowest (pre-pandemic) to the second lowest during the first year of the pandemic. Many potential causes exist for geographic differences in adult obesity rates, including spatial differences in the food and physical activity environments. For example, Wende et al. (2021) showed that the South had significantly lower food and physical activity environment scores than other regions. Michimi and Wimberly (2015) reported a higher concentration of convenience store and fast-food workers in the South and Midwest. Higher concentrations of convenience store and fast-food workers are associated with higher community obesity rates compared to communities with larger concentrations of supermarket and full-service restaurant workers.

Conclusion

During the first year of the COVID-19 pandemic in the United States, obesity rates increased across the overall adult population, but there were not significant changes among every adult subgroup examined. Some subgroups experienced substantive increases in obesity growth rates, especially younger adults, low-education adults, those in low-income SNAP-eligible households, and adults living in the West Census region.

Previous reports showed that unemployment is typically associated with higher obesity rates (e.g., Biener et al., 2018; Cawley, 2004), but the authors found that obesity among the unemployed was unchanged during the first year of the pandemic. However, the small increase in obesity among employed individuals corresponded with the simulated predictions of Chen et al. (2018). They reported that reemployment could lead to an increase in the prevalence of population-wide obesity through a reduction in leisure time. Geographic trends also deviated from pre-pandemic trajectories. The West Census region experienced the greatest increase in obesity. These findings contrasted with long-term trends of greater obesity rate increases in rural areas of the United States and the South (Befort et al., 2012; Lundeen et al., 2018). Finally, although middle-aged adults had the highest rate of obesity before the pandemic (Hales et al., 2020b), young adults reported the highest obesity growth rates during the pandemic. This finding is consistent with that of Freedman et al. (2022), who found that young adults gained more weight than older adults during the pandemic. The pandemic disruption shifted increased weight accumulation from middle-aged to younger adults during the early months of the pandemic.

Other intra-pandemic obesity patterns were more closely predicted by pre-pandemic trends. Findings within racial and ethnic categories suggest that pre-pandemic obesity rate disparities were perpetuated during the first year of the pandemic. Like earlier trends (e.g., Hicken et al., 2018), non-Hispanic Black adults were more susceptible to obesity from 2019 to 2020. The higher obesity rate among women was consistent with pandemic-

era findings that women gained more weight than men during the first year of the pandemic (Freedman et al., 2022). Findings were also consistent with prior reports that the presence of children in the household during the pandemic decreased physical activity, especially among women (e.g., Matthews et al., 2022), and exposed adults to greater obesity risk. The increased obesity rates among low-income adults were accordant with previous research identifying low income as a possible obesity-related risk factor (Grecu and Rotthoff, 2015). Finally, the inverse relationship between educational attainment and obesity growth rates during the pandemic was consistent with pre-pandemic findings from developed economies (Cohen et al., 2013).

Groups with the most limited resources were less likely to contain obesity rate growth during the first year of the pandemic. Younger adults and those with limited income and education were most at risk of increased obesity.

The research was limited to the demographic and socioeconomic characteristics available in the BRFSS dataset, and data measuring food-related mechanisms were unavailable. The authors were also limited to comparisons between categorically defined groups. The analysis did not have groups defined by multiple categories to allow for intersectional comparisons (e.g., low-income females compared to high-income female obesity growth rates). Due to the lack of data on food-related mechanisms during the pandemic, the authors were not able to explore whether and how much increases in obesity were fueled by changes in access to healthy foods or shifts in demand toward calorie-dense foods. Evidence suggests food spending changed by food retail type (e.g., grocery retail versus foodservice retail) and across food categories during the first year of the pandemic (Okrent and Zeballos, 2022). For example, steep declines in food-away-from-home spending were mostly offset by food-at-home spending, with protein foods, desserts, prepared meals, and snacks contributing most to the increase in the food-at-home budget share (Okrent and Zeballos, 2022). However, the authors were unable to link actual food consumption to weight outcomes because pandemic restrictions did not permit in-person dietary interviews or anthropometric measurements that would have otherwise been collected as part of the National Health and Nutrition Examination Survey (NHANES) (Centers for Disease Control and Prevention, 2022b). There were also additional society- and economy-wide shifts during the first year of the COVID-19 pandemic. These included expanded unemployment insurance or food distribution disruptions that were not considered in the analysis. Despite these limitations, this study identifies dynamic changes in obesity outcomes across population sub-groups during the first year of the pandemic. The information about subgroup differences may inform differentiated analysis moving forward.

References

- Almada, L.N., and R. Tchernis. 2018. "Measuring Effects of SNAP on Obesity at the Intensive Margin," *Economics & Human Biology* 31:150–163.
- Almandoz, J., L. Xie, J.N. Schellinger, M.S. Mathew, E. Morales Marroquin, N. Murvelashvili, S. Khatiwada, S. Kukreja, C. McAdams, and S.E. Messiah. 2022. "Changes in Body Weight, Health Behaviors, and Mental Health in Adults with Obesity During the COVID-19 Pandemic," *Obesity* 30(9):1875–1886.
- Befort, C.A., N. Nazir, and M.G. Perri. 2012. "Prevalence of Obesity Among Adults From Rural and Urban Areas of the United States: Findings From NHANES (2005-2008)," *The Journal of Rural Health* 28(4):392–397.
- Belanger, M.J., M.A. Hill, A.M. Angelidi, M. Dalamaga, J.R. Sowers, and C.S. Mantzoros. 2020. "COVID-19 and Disparities in Nutrition and Obesity," *New England Journal of Medicine* 383(11):e69.
- Bell, C.N., L.B.T. Carr, M.P. Chaparro, C.T.S. Tobin, and K.P. Theall. 2021. "Household Composition, Income, and Body Mass Index Among Adults by Race/Ethnicity and Sex," *Journal of Racial and Ethnic Health Disparities*: 9(4):1488–1499.
- Biener, A., J. Cawley, and C. Meyerhoefer. 2018. "The Impact of Obesity on Medical Care Costs and Labor Market Outcomes in the US," *Clinical Chemistry* 64(1):108–117.
- Cawley, J. 2004. "The Impact of Obesity on Wages," *Journal of Human Resources* 39(2):451–474.
- Cawley, J., J.C. Maclean, M. Hammer, and N. Wintfeld. 2015. "Reporting Error in Weight and its Implications for Bias in Economic Models," *Economics of Human Biology* 19:27–44.
- Centers for Disease Control and Prevention. 2009. "Study Estimates Medical Cost of Obesity May Be as High as \$147 Billion Annually," U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, Atlanta, Georgia.
- Centers for Disease Control and Prevention. 2013. "The BRFSS Data User Guide," U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, Division of Population Health, Atlanta, Georgia.
- Centers for Disease Control and Prevention. 2022a. "CDC Orders," U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, Atlanta, Georgia (accessed June 14, 2022).
- Centers for Disease Control and Prevention. 2022b. "People with Certain Medical Conditions," U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, Atlanta, Georgia (accessed June 14, 2022).
- Centers for Disease Control and Prevention. 2022c. "Defining Adult Overweight and Obesity," U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, Atlanta, Georgia.
- Centers for Disease Control and Prevention. 2022d. "NHANES Questionnaires, Datasets, and Related Documentation," U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, Atlanta, Georgia.

- Chen, H.J., H. Xue, S. Liu, T.T.K. Huang, Y.C. Wang, and Y. Wang. 2018. "Obesity Trend in the United States and Economic Intervention Options to Change It: A Simulation Study Linking Ecological Epidemiology and System Dynamics Modeling," *Public Health* 161:20–28.
- Cohen, A.K., M. Rai, D.H. Rehkopf, and B. Abrams. 2013. "Educational Attainment and Obesity: A Systematic Review," *Obesity Reviews* 14(12):989–1005.
- Coronavirus Resource Center. 2022a. "Number of Daily Deaths," Johns Hopkins University and Medicine, Baltimore, Maryland.
- Coronavirus Resource Center. 2022b. "Weekly Hospitalization Trends," Johns Hopkins University, Baltimore, Maryland.
- Courtemanche, C., J.C. Pinkston, and J. Steward. 2015. "Adjusting Body Mass for Measurement Error with Invalid Validation Data," *Economics of Human Biology* 19:275–293.
- Flanagan, E.W., R.A. Beyl, N. Fearnback, A.D. Altazan, C.K. Martin, and L.M. Redman. 2020. "The Impact of COVID-19 Stay-At-Home Orders on Health Behavior in Adults," *Obesity* 29(2):438–445.
- Freedman, D.S., L. Kompaniyets, C. Daymont, L. Zhao, and H.M. Blanck. 2022. "Weight Gain Among U.S. Adults During the COVID-19 Pandemic through May 2021," *Obesity* 30(10):2064–2070.
- Fryar, C.D., M.D. Carroll, and J. Afful. 2020. *Prevalence of Overweight, Obesity, and Severe Obesity Among Adults Aged 20 and Over: United States, 1960–1962 through 2017–2018*. U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Health Statistics, Hyattsville, Maryland.
- Greco, A.M., and K.W. Rotthoff. 2015. "Economic Growth and Obesity: Findings of an Obesity Kuznets Curve," *Applied Economic Letters* 22(7):539–543.
- Hales, C.M., M.D. Carroll, C.D. Fryar, and C.L. Ogden. 2020. *Prevalence of Obesity and Severe Obesity Among Adults: United States, 2017–2018*. NCHS Data Brief 360, U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Health Statistics, Hyattsville, Maryland.
- Han, T.S., A. Tajar, and M.E. Lean. 2011. "Obesity and Weight Management in the Elderly," *British Medical Bulletin* 97:169–96.
- Hicken, M.T., H. Lee, and A.K. Hing. 2018. "The Weight of Racism: Vigilance and Racial Inequalities in Weight-related Measures," *Social Science & Medicine* 199:157–166.
- Jackson, J.E., M.P. Doescher, A.F. Jerant, and L.G. Hart. 2005. "A National Study of Obesity Prevalence and Trends by Type of Rural County," *The Journal of Rural Health* 21(2):140–148.
- Kirwan, R., D. McCullough, T. Butler, F. Perez de Heredia, I.G. Davies, and C. Stewart. 2020. "Sarcopenia During COVID-19 Lockdown Restrictions: Long-term Health Effects of Short-term Muscle Loss," *GeroScience* 42(6):1547–1578.
- Koh, K., S.C. Grady, J.T. Darden, and I. Vojnovic. 2018. "Adult Obesity Prevalence at the County Level in the United States, 2000–2010: Downscaling Public Health Survey Data Using a Spatial Microsimulation Approach," *Spatial and Spatio-Temporal Epidemiology* 26:153–164.
- Li, Y., T. Cai, H. Wang, and G. Guo. 2021. "Achieved Educational Attainment, Inherited Genetic Endowment for Education, and Obesity," *Biodemography and Social Biology* 66(2):132–144.

- Lundeen, E.A., S. Park, L. Pan, T. O’Toole, K. Matthews, and H.M. Blank. 2018. “Obesity Prevalence Among Adults Living in Metropolitan and Nonmetropolitan Counties—United States, 2016,” *Morbidity and Mortality Weekly Report* 67(653–658).
- Matthews, C.E., P. Saint-Maurice, J.E. Fulton, S. Patel, E. Loftfield, J. Sampson, S.K. Keadle, and D. Berrigan. 2022. “Changes in Physical Activity and Sedentary Time in United States Adults in Response to COVID-19,” *PLOS One* 17(9):e0273919.
- McMillian, M.M., and R.J. Thorpe, Jr. 2021. “Relationships of Educational Attainment and Household Food Insecurity with Obesity: Findings from the 2007–2016 National Health and Nutrition Examination Survey,” *International Journal of Environmental Research and Public Health* 18(15).
- Medvedyuk, S., A. Ali, and D. Raphael. 2018. “Ideology, Obesity, and the Social Determinants of Health: A Critical Analysis of the Obesity and Health Relationship,” *Critical Public Health* 28(5):573–585.
- Michimi, A., and M.C. Wimberly. 2015. “The Food Environment and Adult Obesity in U.S. Metropolitan Areas,” *Geospatial Health* 10(2):223–231.
- National Center for Chronic Disease Prevention and Health Promotion. 2022. “Behavioral Risk Factor Surveillance System.” U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, Atlanta, Georgia.
- National Center for Chronic Disease Prevention and Health Promotion. 2022. “Consequences of Obesity.” U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, Atlanta, Georgia.
- Ogden, C. L. 2009. “Disparities in Obesity Prevalence in the United States: Black Women at Risk,” *American Journal of Clinical Nutrition* 89(4):1001–2.
- Ogden, C.L., T.H. Fakhouri, M.D. Carroll, C.M. Hales, C.D. Fryaer, X. Li, and D.S. Freedman. 2017. “Prevalence of Obesity Among Adults, by Household Income and Education – United States, 2011–2014,” *Morbidity and Mortality Weekly Report* 66(50):1369–1373.
- Okrent, A., and E. Zeballos. 2022. *COVID-19 Working Paper: Consumer Food Spending Changes During the COVID-19 Pandemic*, AP-110, U.S. Department of Agriculture, Economic Research Service, Washington, D.C.
- Onyeaka, H., C.K. Anumudu, Z.T. Al-Sharify, E. Egele-Godswill, and P. Mbaegbu. 2021. “COVID-19 Pandemic: A Review of the Global Lockdown and its Far-reaching Effects,” *Scientific Progress* 104(2):1–18.
- Pierrannunzi, C., M. Town, W. Garvin, F.E. Shaw, and L. Balluz. 2012. “Methodologic Changes in the Behavioral Risk Factor Surveillance System in 2011 and Potential Effects on Prevalence Estimates,” *Morbidity and Mortality Weekly Report* 61(22):410–413.
- Poobalan, A., and L. Aucott. 2016. “Obesity Among Young Adults in Developing Countries: A Systematic Overview,” *Current Obesity Reports* 5(1):2–13.
- Pou, S.A., M.D.P. Diaz, G.A. Velázquez, and L.R. Aballay. 2021. “Sociodemographic Disparities and Contextual Factors in Obesity: Updated Evidence From a National Survey of Risk Factors for Chronic Diseases,” *Public Health Nutrition*: 1–13.
- Restrepo, B. 2022. “Obesity Prevalence Among U.S. Adults During the COVID-19 Pandemic,” *American Journal of Preventive Medicine* 63:1.
- Sanchis-Gomar, F., C.J. Lavie, M.R. Mehra, B.M. Henry, and G. Lippi. 2020. “Obesity and Outcomes in COVID-19: When an Epidemic and Pandemic Collide,” *Mayo Clinic Proceedings* 95(7):1445–1453.

- Singh, G. K., J. N. DiBari, and H. Lee. 2021. “Prevalence and Social and Built Environmental Determinants of Maternal Pre-pregnancy Obesity in 68 Major Metropolitan Cities of the United States, 2013–2016,” *Journal of Environmental and Public Health* 2021:16.
- Sobal, J., B.S. Rauschenbach, and E.A. Frongillo. 1992. “Marital Status, Fatness and Obesity,” *Social Science & Medicine* 35(7):915–923.
- Solar, O, and A. Irwin. 2010. *A Conceptual Framework for Action on the Social Determinants of Health*. World Health Organization, Geneva, Switzerland.
- Stefan, N., A.L. Birkenfeld, and M.B. Schulze. 2021. “Global Pandemics Interconnected—Obesity, Impaired Metabolic Health and COVID-19,” *Nature Reviews Endocrinology* 17(3):135–149.
- The Economist. 2020. “COVID-19 Seems to Have Changed Lifestyles for Good,” *The Economist*, August 8.
- The White House. 2022. Biden-Harris Administration National Strategy on Hunger, Nutrition, and Health. Washington, D.C.
- Thompson, S.A., Y. Serkez, and L. Kelley. 2020. “How Has Your State Reacted to Social Distancing?” *The New York Times*, March 23.
- U.S. Bureau of Labor Statistics. 2020. “The Employment Situation—March 2020,” U.S. Department of Labor, Washington, DC.
- U.S. Department of Agriculture. 2022. “USDA Actions on Nutrition Security,” United States Department of Agriculture, Washington, DC.
- U.S. Department of Commerce, Bureau of the Census. 2022. “West Region is Made up of 13 States,” United States Department of Commerce, Washington, DC.
- Ver Ploeg, M., L. Mancino, and B. Lin. 2007. *Food and Nutrition Assistance Programs and Obesity: 1976–2002*, ERR-48, U. S. Department of Agriculture, Economic Research Service. Washington, D.C.
- Ver Ploeg, M., L. Mancino, B. Lin, and C. Wang. 2007. “The Vanishing Weight Gap: Trends in Obesity Among Adult Food Stamp Participants (U.S.) (1976–2002),” *Economics and Human Biology* 5(1):20–36.
- Wang, Y., M.A. Beydoun, J. Min, H. Xue, L.A. Kaminsky, and L.J. Cheskin. 2020. “Has the Prevalence of Overweight, Obesity and Central Obesity Levelled Off in the United States? Trends, Patterns, Disparities, and Future Projections for the Obesity Epidemic,” *International Journal of Epidemiology* 49(3):810–823.
- Wells, J.C.K., A.A. Marphatia, T.J. Cole, and D. McCoy. 2012. “Associations of Economic and Gender Inequality with Global Obesity Prevalence: Understanding the Female Excess,” *Social Science & Medicine* 75(3):482–490.
- Wende, M.E., E.W. Stowe, J.M. Eberth, A.C. McLain, A.D. Liese, C.B. Breneman, M.J. Josey, S. Morgan Hughey, and A.T. Kaczynski. 2021. “Spatial Clustering Patterns and Regional Variations for Food and Physical Activity Environments Across the United States,” *International Journal of Environmental Health Research* 31(8):976–990.
- World Health Organization. 2022. “Obesity and Overweight,” World Health Organization, Geneva, Switzerland.